

**Results of the Czechoslovak-Iranian entomological expeditions to Iran.
Coleoptera: Silphidae**

**Výsledky československo-iránských entomologických expedicí do Íránu.
Coleoptera: Silphidae**

Jan RŮŽIČKA

Department of Ecology, Faculty of Forestry, Czech Agricultural University, CZ-165 21 Praha 6,
Czech Republic

Faunistics, Coleoptera, Silphidae, Iran

Abstract. The faunistic data are given for 68 specimens, representing seven species of Silphidae, collected during 1970-1977 by the Czechoslovak-Iranian entomological expeditions in Iran. Recent finding of *Aclypea cicatricosa* Reitter, 1885 and first reliable records for *Nicrophorus vespilloides* Herbst, 1784 and *N. investigator* Zetterstedt, 1824 from Iran are provided.

Three expeditions to Iran were undertaken by the Department of Entomology of the National Museum (Natural History) in Prague in 1970, 1973 and 1977, respectively. The basic data about the expeditions, including detailed description of collecting localities (numbering successively), maps and photos can be find in Hoberlandt (1974, 1981, 1983).

In this paper, the material of carrion beetles (Coleoptera: Silphidae) is elaborated. All the material is deposited in collections of National Museum, Praha, unless otherwise mentioned.

***Thanatophilus sinuatus* (Fabricius, 1775)**

Material examined. N. Iran, Tehran prov., central Elburz Mts, Kandavan - pass, 2700-2900 m, south slope, loc. no. 395, 4-9.vii.1977, 1 ♂.

Palearctic species, from Iran reported by Schawaller (1981).

***Ablattaria cribrata* (Ménétriés, 1832)**

Material examined. N. Iran, Gilan, Sheykh Mahalleh, 160 m, loc. no. 390, 28.vi.-3.vii.1977, 1 ♀. This species is known only from southern parts of Russia and the Caucasus, a single report from northwestern Iran is reported by Schawaller (1979).

***Aclypea undata* (Müller, 1776)**

Material examined. N. Iran: Mazandaran, central Elburz Mts, Damavand, Lajran, 2400 m, loc. no. 68, 23.vii.1970, 1 ♂ 2 ♀♀; Tehran prov., Elburz Mts Central, Kandavan, Val. N of tunnel, 2545 m, loc. no. 86, 10-11.vi.1970, 1 ♂; Tehran prov., Elburz Mts Central, Kandavan - pass, 2700-2900 m, south slope, loc. no. 395, 4-9.vii.1977, 1 ♀; Tehran prov., 8 km W of Gachsar, loc. no. 396, 6.-8.vii.1977, 1 ♀; Elburz Mts, Damavand Mt., vii.1976, J. Boháč lgt., J. Růžička coll., 1 ♀. Palearctic species, the specimens from the Caucasus, the Transcaucasia, Syria and "Persia" with more lustrous and bumpy elytra were described as ssp. *verrucosa* (Ménétriés, 1832) - Reitter (1885), Portevin (1926) and Nikolaev (1990); it is recently considered as being a junior synonym of *A. undata* by Schawaller (pers. comm.).

***Aclypea cicatricosa* Reitter, 1885**

Material examined. E. Iran, Kerman prov., Jebal Barez Mts, Deh Bakri, 1700-1750 m, loc. no. 186, 30.iv.-3.v.1973, 6 ♂♂ 14 ♀♀, some specimens also deposited in the author's collection and in coll. Schawaller.

The species was described after a single female specimen from "Persia" (Reitter 1885), three further records from western Turkmenia have been recently provided by Nikolaev (1990).

***Silpha obscura obscura* Linnaeus, 1758**

Material examined. N.W. Iran, Azarbaijan, Sufian, 30 km W Tabriz, loc. no. 27, 20-21.vi.1970, 3 ♂♂ 1 ♀; N.W. Iran, Tehran prov., Ghazvin, loc. no. 29, 24.vi.1970, 2 ♂♂ 2 ♀♀; N. Iran, Mazandaran, Elburz Mts Central, Gazanak, in the valley of the river Haraz Chay, 1400 m, loc. no. 63, 20-21.vii.1970, 8 ♂♂ 4 ♀♀; N. Iran, Tehran prov., central Elburz Mts, Kandavan, valley N of tunnel, 2545 m, loc. no. 86, 1 ♂ 2 ♀♀.

Nominotypical subspecies of this palaearctic species, reported from Iran by Schawaller (1980).

***Silpha obscura nitida* Portevin, 1907**

Material examined. N. Iran, Tehran prov., Elburz Mts West, Rudbarak, northern slope of the Alam Kuh massif, plain of the Kalardasht, 1850-2400 m, loc. no. 92, 13.viii.1970, 4 ♂♂ 3 ♀♀; N. Iran, Gilan, Talesh Mts, 24 km N of Khalkhal, Pushte-Kuh, pass, 2200-2300 m, loc. no. 95, 13.viii.1970, 4 ♂♂ 5 ♀♀.

This subspecies is restricted to the Caucasus, the Transcaucasia, northeastern Turkey; from north-western and northern Iran reported by Schawaller (1980).

***Nicrophorus vespilloides* Herbst, 1784**

Material examined. N. Iran, Tehran prov., Elburz Mts West, Rudbarak, northern slope of the Alam Kuh massif, plain of the Kalardasht, 1850-2400 m, loc. no. 92, 13.viii.1970, 1 ♀.

Holarctic species, the first reliable record from Iran.

***Nicrophorus investigator* Zetterstedt, 1824**

Material examined. N. Iran, Tehran prov., Elburz Mts West, Rudbarak, northern slope of the Alam Kuh massif, plain of the Kalardasht, 1850-2400 m, loc. no. 92, 13.viii.1970, 1 ♀.

Holarctic species, the first reliable record from Iran.

ACKNOWLEDGEMENTS. I am obliged to Dr J. Jelínek for the possibility to study the material from the expeditions of National Museum, Praha, to Iran. My thanks are due to Dr W. Schawaller, Stuttgart, for the revision of the identity of *Aclypea cicatricosa* and *Ablattaria cribrata*.

REFERENCES

- HOBERLANDT L. 1974: Results of the Czechoslovak-Iranian entomological expedition to Iran 1970. No. 1: Introduction. *Acta Entomol. Mus. Nat. Pragae, Suppl.* 6: 9-20.
- HOBERLANDT L. 1981: Results of the Czechoslovak-Iranian entomological expeditions to Iran. Introduction to the Second expedition 1973. *Acta Entomol. Mus. Nat. Pragae*, 40: 5-32.
- HOBERLANDT L. 1983: Results of the Czechoslovak-Iranian entomological expeditions to Iran. Introduction to the Third expedition 1977. *Acta Entomol. Mus. Nat. Pragae*, 41: 5-24.
- NIKOLAEV G. V. 1990: [A review of the genus *Aclypea* Rtt. (Coleoptera, Silphidae) from the USSR]. *Trudy Inst. Zool. Akad. Nauk Kazakhskoi SSR*, 45: 38-45 (in Russian).
- PORTEVIN G. 1926: *Les grands nécrophages du globe, Silphini - Necrodini - Necrophorini*. Paul Lechevalier, Paris, 270 pp.
- REITTER E. 1885: Bestimmungs-Tabellen der europäischen Coleopteren 12. Necrophaga (Platysyllidae, Leptinidae, Silphidae, Anisotomidae und Clambidae). *Verh. Naturforsch. Ver. Brünn*, 23: 1-122.
- SCHAWALLER W. 1979: Revision der Gattung *Ablattaria* Reitter 1884 (Coleoptera: Silphidae). *Stuttgarter Beitr. Naturk. (A)*, 321: 1-8.
- SCHAWALLER W. 1980: *Silpha obscura*, ein Beispiel für Subspecies-Differenzierung bei Käfern (Coleoptera, Silphidae). *Stuttgarter Beitr. Naturk. (A)*, 334: 1-11.
- SCHAWALLER W. 1981: Die Aaskäfer des Himalaya (Insecta: Coleoptera: Silphidae s. str.). *Senckenbergiana Biol.*, 62: 237-260.

SOUHRN

V práci je zpracován materiál mrchožroutovitých brouků čeledi Silphidae, sebraný během let 1970-1977 účastníky tří československo-iránských entomologických expedicí do Íránu. Celkem byl zpracován materiál 68 exemplářů. Ze sedmi druhů mrchožroutů je nejzajímavější nález druhu *Aclypea cicatricosa*, popsaného Reitterem (1885) podle jediné samice z "Persie" (bez bližší lokalizace), druh je jinak známý pouze ze západní Turmenie (Nikolaev 1990). První konkrétní nálezy z Íránu jsou také uvedeny pro holarktické druhy hrobaříků *Nicrophorus vespilloides* Herbst a *N. investigator* Zetterstedt.